

# Inter- and intra-specific variation in hair cortisol concentrations of Neotropical bats

Natalia I. Sandoval-Herrera<sup>1,2,\*</sup>, Gabriela F. Mastromonaco<sup>3</sup>, Daniel J. Becker<sup>4</sup>, Nancy B. Simmons<sup>5</sup> and Kenneth C. Welch Jr<sup>1,2</sup>

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of Toronto, Ontario, M5S 3B2, Canada

<sup>2</sup>Department of Biological Sciences, University of Toronto Scarborough, Ontario, M1C 1A4, Canada

<sup>3</sup>Reproductive Sciences, Toronto Zoo, Ontario, M1B 5K7, Canada

<sup>4</sup>Department of Biology, University of Oklahoma, Norman, OK 73019, USA

<sup>5</sup>Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural History, New York, NY, 10024-5102, USA

\*Corresponding author: Department of Ecology and Evolutionary Biology, University of Toronto, Ontario, Canada.  
Email: natalia.sandovalherrera@mail.utoronto.ca

Quantifying hair cortisol has become popular in wildlife ecology for its practical advantages for evaluating stress. Before hair cortisol levels can be reliably interpreted, however, it is key to first understand the intrinsic factors explaining intra- and inter-specific variation. Bats are an ecologically diverse group of mammals that allow studying such variation. Given that many bat species are threatened or have declining populations in parts of their range, minimally invasive tools for monitoring colony health and identifying cryptic stressors are needed to efficiently direct conservation efforts. Here we describe intra- and inter-specific sources of variation in hair cortisol levels in 18 Neotropical bat species from Belize and Mexico. We found that fecundity is an important ecological trait explaining inter-specific variation in bat hair cortisol. Other ecological variables such as colony size, roost durability and basal metabolic rate did not explain hair cortisol variation among species. At the individual level, females exhibited higher hair cortisol levels than males and the effect of body mass varied among species. Overall, our findings help validate and accurately apply hair cortisol as a monitoring tool in free-ranging bats.

**Editor:** Steven Cooke

Received 20 February 2021; Revised 13 June 2021; Editorial Decision 13 June 2021; Accepted 8 July 2021

**Cite as:** Sandoval-Herrera NI, Mastromonaco GF, Becker DJ, Simmons NB, Welch KC (2021) Inter- and intra-specific variation in hair cortisol concentrations of Neotropical bats. *Conserv Physiol* 9(1): coab053; doi:10.1093/conphys/coab053.

## Introduction

Free-living animals face multiple natural and anthropogenic challenges that threaten their survival and thus are of considerable interest to ecophysiologicals concerned with the study of effects of stress on vertebrates. One of the most extensively studied processes associated with response to stressors (biotic or abiotic environmental factors that disrupt homeostasis; Schulte, 2014) is the release of glucocorticoid (GC) hormones (Creagh and Brendan Delehanty, 2013; MacDougall-Shackle-

ton *et al.*, 2019). GCs are known to facilitate the mobilization of energy required to cope with stressors and, during normal conditions, play a key role in regulating growth, circadian activity and energy metabolism (review in Landys *et al.*, 2006). Levels of GCs are commonly employed as a biomarker of allostatic load or stress (indirect indicators of health) (Sapolsky *et al.*, 2000; Wikelski and Cooke, 2006; Pearson Murphy, 2007; Busch and Hayward, 2009). GC secretion is a well-conserved process across vertebrates and involves activation of the hypothalamic–pituitary–adrenal (HPA) axis

and release of GCs from the adrenal glands to the blood stream (Norris and Carr, 2013). In mammals, the primary GC is cortisol, which induces a cascade of events to maintain homeostasis at multiple target tissues (Pearson Murphy, 2007; Boonstra, 2013). An acute increase in GC levels can benefit an individual's survival (e.g. by allocating energy in defence and escape) yet if adverse conditions remain, continuously elevated GCs in circulation can become pathological, causing immune suppression, neuronal cell death and reproductive impairment (Sapolsky *et al.*, 2000; Tilbrook, 2000; Wingfield and Romero, 2011; Hing *et al.*, 2016).

Although many of the environmental challenges that wild populations experience are chronic (e.g. prolonged food deprivation, climate change, habitat disturbance, pollution), studies of stress physiology have focused on detecting acute stress by looking at GC levels in blood, urine and faeces (Sheriff *et al.*, 2011; Creagh and Brendan Delehanty, 2013). The rapid turnover of these tissues, however, only gives short-term information of HPA activity over periods of hours or days (Sheriff *et al.*, 2011), which may not be an appropriate time scale. Assessment of cortisol in tissues with slower turnover rates, such as hair, may reflect circulating cortisol levels over longer periods of several weeks or even months, which is the time scale over which chronic environmentally induced stress would be expected to occur (Davenport *et al.*, 2006; Macbeth *et al.*, 2010; Ashley *et al.*, 2011; Mastromonaco *et al.*, 2014). Cortisol is incorporated into developing hairs from the blood stream during periods of active hair growth, allowing researchers to retrospectively examine cortisol production at the time that a stressor or stressors were faced (Davenport *et al.*, 2006; Pragst and Balikova, 2006). Hair can be collected in a minimally invasive manner, is usually easily accessible in relatively large amounts and is easy to store and transport, all of which make it particularly useful for wildlife studies, especially those involving threatened or endangered species (Koren *et al.*, 2002; Macbeth *et al.*, 2010, 2012). Hair cortisol levels are not likely affected by stress induced by capture and/or handling, which is one of the main limitations of blood GC analysis (Russell *et al.*, 2012). A single sample of hair can also provide complementary and valuable information about ecology and behaviour, including diet and movement (e.g. using stable isotope analyses; Fraser *et al.*, 2010; Sullivan *et al.*, 2012; Voigt *et al.*, 2012; Oelbaum *et al.*, 2019), condition (e.g. nutrition; Montillo *et al.*, 2019), toxicant exposure (Hernout *et al.*, 2016; Becker *et al.*, 2018) and molecular identification (Magioli *et al.*, 2019), opening possibilities for more integrative studies. However, analyses of hair samples can be challenging. Despite being a very promising tool for assessing wildlife health, quantifying hair cortisol is a method that has limitations; although these are largely based on lack of detailed knowledge of patterns of hair growth (Meyer and Novak, 2012; Russell *et al.*, 2012; Sharpley *et al.*, 2012). For example, the exact time scale reflected in any given sample will depend on the rate of hair growth and moulting patterns; this information is unknown for most species, which makes

the time window being evaluated unclear (Koren *et al.*, 2002; Fourie *et al.*, 2016). Moreover, rates of cortisol incorporation to the hair shaft are known to differ across body regions and among species (Sharpley *et al.*, 2012; Acker *et al.*, 2018; Lavergne *et al.*, 2020). Nevertheless, hair cortisol levels offer a potentially powerful tool for assessing relatively long-term stress levels in mammals.

Hair cortisol and its correlation with natural and anthropogenic stressors has been explored for different wild mammals, including rhesus monkeys (*Macaca mulatta*; Dettmer *et al.*, 2014), grizzly bears (*Ursus arctos*; Macbeth *et al.*, 2010), reindeer/caribou (*Rangifer tarandus*; Ashley *et al.*, 2011), lynx (*Lynx canadensis*; Terwissen *et al.*, 2013; Azevedo *et al.*, 2020), mongoose (*Herpestes ichneumon*; Azevedo *et al.*, 2019) and snowshoe hares (*Lepus americanus*; Lavergne *et al.*, 2020); other examples reviewed by Kalliokoski *et al.*, 2019). Although most of these studies support hair cortisol as an informative measure of central HPA activity, they also identified intrinsic factors such as age, sex, reproductive stage and social status that modulate GC levels in different contexts (Wingfield and Romero, 2011; Crespi *et al.*, 2013; Hau *et al.*, 2016). Not accounting for these intrinsic sources of variation in GC levels may lead to incorrect or misleading estimates of the effects of stressors on individual fitness and population health (Sapolsky *et al.*, 2000; Reeder and Kramer, 2005; Busch and Hayward, 2009; Wingfield and Romero, 2011; Kalliokoski *et al.*, 2019).

Ecological traits such as diet, fecundity and lifespan, as well as phylogenetic relatedness, have been proposed to explain differences in baseline cortisol levels in wild species (Wingfield and Romero, 2011; Patterson *et al.*, 2014). Evolution of different life-history strategies are also thought to have led to different adaptations in HPA activity modulation so as to maximize individual fitness within species (Bonier *et al.*, 2009; Bonier and Martin, 2016). Bats are a very ecologically diverse group comprising over 1400 species that live in most terrestrial ecosystems and have a wide variety of diets, use many different roost types and have many different social systems (Kunz and Fenton, 2005; Dumont *et al.*, 2012; Gunnell and Simmons, 2012; Simmons and Cirranello, 2020). This diversity provides the opportunity to study the ecological correlates of cortisol levels among phylogenetically related species with different life history traits. Few ecological correlates of GCs have been evaluated simultaneously in mammalian groups in the context of cortisol studies, and fewer studies have further related cortisol levels to life history traits across multiple species from a single mammalian clade. Among bats, variation in plasma cortisol levels associated with seasonal food availability has been studied in two species with contrasting diets, *Carollia perspicillata* and *Desmodus rotundus* (Lewanzik *et al.*, 2012), but no other comparative studies have been conducted within this order. Furthermore, little is known about the modulation of the stress response in bats, despite Chiroptera being the second-most speciose order of mammals.

Bat populations are declining worldwide due to ongoing habitat destruction and land use changes, increased interaction with human environments and associated threats including wind turbine fatalities, hunting and targeted killing, pesticide exposure and emerging infectious diseases such as white-nose syndrome (O'Donnell, 2000; Mickleburgh *et al.*, 2002; Kunz *et al.*, 2007; Frick *et al.*, 2010; Racey, 2013; Voigt and Kingston, 2015). Because many bat species are threatened or have declining populations in parts of their range (IUCN Red List of Threatened Species, 2020), minimally invasive tools to monitor colony health and identify cryptic stressors are critically needed to efficiently direct conservation efforts. It is essential to investigate the factors influencing baseline GCs to properly detect elevated cortisol levels due to long-term stressors.

In this study, we describe intra- and inter-specific sources of variation in baseline hair cortisol levels in bats, which contributes to better understanding the potential for hair cortisol to be an indicator of HPA activity in this taxon. We hypothesize that interspecific variation in hair cortisol of bats will be greater than intra-specific variation and that such heterogeneity will be best explained by ecological traits directly related to energy expenditure, such as basal metabolic rate (BMR), dietary guild, foraging style and roost durability. We expect that species with high energetic demands or less predictable energy acquisition (e.g. less reliable food sources) will have higher hair cortisol. Specifically, we predict the following: (i) a positive relationship between BMR and hair cortisol; (ii) bats that feed on fruit and nectar, which are energy rich and readily available, will have lower hair cortisol; (iii) bats that actively hunt prey during flight, such aerial hawkers, will have higher GC levels owing to greater energetic demands compared to gleaners that can hunt from perches (Norberg and Rayner, 1987; Fenton, 1990); and (iv) species using more ephemeral day roosts (e.g. foliage or crevices under exfoliating bark) will have higher hair cortisol than species using more stable structures (Kunz and Fenton, 2005).

## Material and methods

### Study sites

We sampled bats from northern Belize (Orange Walk District) and two locations in Mexico (Colima and Chihuahua States). In each region, we sampled sites with different levels of habitat fragmentation and agricultural intensity. We used the global Human Modification Index (HMI; Kennedy *et al.*, 2019) as a standardized measure of disturbance, using a 5-km buffer around each collection site. The HMI is a cumulative measurement with possible values between 0 (no disturbance) and 1 (highest disturbance) that includes transportation, human settlement, agriculture, extractive activities and electric infrastructure (Kennedy *et al.*, 2019). Sites were classified as low ( $0 \leq \text{HMI} \leq 0.10$ ), moderate ( $0.10 < \text{HMI} \leq 0.40$ ), high ( $0.40 < \text{HMI} \leq 0.70$ ) and very high ( $0.70 < \text{median HMI} \leq 1.00$ ). At all sites, bats were


captured from 18:00 to 22:00 using mist nets and from 18:00 to 5:00 using harp traps (only in Belize) set along flight paths. Bats sampled during the day were captured in their roosts, mainly caves, using hand nets. We recorded sex, size (body mass, g) and reproductive stage (females: pregnant, lactating; males: active, inactive; Kunz and Parsons, 2009). Reproductive stage was assessed by checking for the presence of scrotal testes in males (indicating that the individual was reproductively active at the time of capture) and by the evidence of pregnancy or lactation (enlarge nipples) in females (Racey, 2009). Only adult individuals were sampled for hair collection.

In Colima (west central Mexico) in March 2019 (dry season), we sampled bats roosting in three caves surrounded by different levels of disturbance: Don Pancho Cave (moderate disturbance, HMI=0.38), El Salitre Cave (high disturbance, HMI=0.44) and La Fábrica Cave (high disturbance, HMI=0.57; Fig. 1). Don Pancho Cave is located on San Agustín island, 1 km away from the coast of Chamela Bay, Jalisco ( $19.5353^{\circ}\text{N}$ ,  $-105.0881^{\circ}\text{W}$ ). El Salitre Cave is near Los Ortices village, Colima ( $19.083330^{\circ}\text{N}$ ,  $-103.726667^{\circ}\text{E}$ ). La Fábrica Cave is 6.4 km southwest of Coquimatlan town, Colima ( $19.1513^{\circ}\text{N}$ ,  $-103.8353^{\circ}\text{W}$ ). We refer here to these locations collectively as central Mexico, and we took samples from three species in this region: *Leptonycteris yerbabuena*, *Macrotus waterhousii*, and *Pteronotus mexicanus*. We also sampled bats foraging close to pecan nut croplands near the town of Jimenez, Chihuahua (northern Mexico). This region is entirely dedicated to the production of pecan nuts with thousands of squared kilometres of cultivated land (Orona Castillo *et al.*, 2018). We visited one that farms using organic practices and another that farms with intensive use of pesticides. However, the estimated HMI was the same for the two sites (HMI=0.49, high disturbance). We collected hair samples from three bat species (*Antrozous pallidus*, *Tadarida brasiliensis* and *Myotis velifer*) at both northern Mexico sites.

Our field sites in Belize consisted of two forest patches of very different size located ~10 km apart and separated by a heterogeneous, largely agricultural landscape. Lamanai Archaeological Reserve (LAR) is a protected secondary semi-deciduous forest of 450 ha with a high canopy and with relatively low disturbance (HMI=0.17) (Herrera *et al.*, 2018). In contrast, the Ka'Kabish archaeological site (KK) is a small remnant forest patch of ~45 ha surrounded by cattle pastures and local croplands (Fig. 1). Although the landscape in Belize is apparently disturbed and highly fragmented, agricultural activity and urban development is not as intense as the field sites in Mexico, which is reflected in their moderate HMI scores (LAR: 0.17; KK: 0.18). We collected hair samples from 12 different species (Table 1) in April 2018 and 2019 (dry season) at Belize sites.

### Ethical statement

Field procedures followed guidelines for safe and humane handling of bats published by of the American Society of



**Figure 1:** Sampling sites in central Belize and Mexico showing the use of land in the surrounding areas. Sources: Sistema Nacional de Información Estadística y Geográfica de Mexico (INEGI, 2013) and Biodiversity and Environmental Resource Data System for Belize.

Mammalogists (Sikes and Bryan, 2016) and were approved by the Institutional Animal Care and Use Committees of the University of Georgia (A2014 04-016-Y3-A5), University of Toronto (20012113) and American Museum of Natural History (AMNHACUC-20180123). Fieldwork was authorized by the Belize Forest Department under permits WL/2/1/18(16) and WL/1/19(06). Sample collection in Mexico was approved under the permit #FAUT-0069.

### Sample collection

We manually trimmed a hair sample (3–10 mg) using round tip curved dissection scissors from the scapular region on the back of each bat. The resulting samples were placed individually in 1–2-ml plastic tubes using flat tweezers. The dissection tools were cleaned with 70% ethanol between sampling different individuals to avoid cross contamination. The amount of hair removed from each bat depended on the hair density of each species. The hair shaft was carefully cut close to the root avoiding removing skin or follicle tissue. From pilot analyses, we determined a minimum amount of 3 mg of hair was necessary to obtain values around 50% binding on the standard curve thereby accurately estimating cortisol concentration in the sample.

### Extraction and quantification of cortisol

Hair samples were processed and analysed at the endocrinology laboratory at the Toronto Zoo following methods described by Acker *et al.*, 2018. Each hair sample was spread apart and weighed in a 7-ml glass scintillation vial. To avoid contamination with other biological fluids, all hair samples were washed with 100% methanol by vortexing in a tube for 10 s and immediately removing the methanol using a pipettor. Immediately thereafter, 100% methanol was added to each sample, at a ratio of 0.005 g/ml. Samples were then mixed for 24 hrs on a plate shaker (MBI Orbital Shaker; Montreal Biotechnologies Inc., Montreal, Quebec City, Canada). After 24 hrs the vials were centrifuged for 10 min at 2400 g. The supernatants were pipetted off into clean glass vials and dried down under air in a fume hood. The dried extracts were stored at  $-20^{\circ}\text{C}$  until analysis.

Samples were brought to room temperature prior to analysis. Reconstitution of the desiccated extracts was done by adding phosphate buffer and vortexing for 10 s. Belize samples were reconstituted neat (i.e. evaporated 150  $\mu\text{l}$  and reconstituted with 150  $\mu\text{l}$ ), and Mexico samples were reconstituted as follows: three species were neat, two species diluted 1:5 and one species diluted 1:50 in phosphate buffer (Andreasson *et al.*, 2015). Cortisol concentrations were determined using an enzyme immunoassay (EIA) previously described

**Table 1:** Species-level ecological traits and hair cortisol data for 18 Neotropical bat species

Family	Species	Region	N	Mass(g)	Dietary guild	Foraging style	WAR	Roost durability	Fecundity (litter/yr)	Colony size	Lifespan (years)	Cortisol (ng/g)	
												Mean	SD
Emballonuridae	<i>R. naso</i>	BZ	14	4.54 ± 1.47	Animalivory	Aerial forager	6.5	2	>1	Small	5	639.57	392.10
												±	±
Molossidae	<i>Saccopterix billineata</i>	BZ	20	6.79 ± 1.12	Animalivory	Aerial forager	6.1	2.4	1	Small	5	347.28	232.01
												±	±
Molossidae	<i>M. nigricans</i>	BZ	5	35.00 ± 2.35	Animalivory	Aerial forager	11.1	4	1	Large	10	227.80	123.54
												±	±
Mormoopidae	<i>T. brasiliensis</i>	MMX	23	10.73 ± 0.63	Animalivory	Aerial forager	8.6	5.3	1	Large	10	1782.87	751.07
												±	±
Mormoopidae	<i>P. mesoamericanus</i>	BZ	26	17.08 ± 2.06	Animalivory	Aerial forager	6.7	5.6	1	Large	10	218.79	76.48
												±	±
Phyllostomidae	<i>P. mexicanus</i>	CMX	35	13.37 ± 1.18	Animalivory	Aerial forager	6.7	5.6	1	Large	10	1246.96	1663.56
												±	±
Phyllostomidae	<i>D. rotundus</i>	BZ	22	28.01 ± 3.54	Animalivory	Gleaner	6.7	4.5	>1	Large	15	395.00	226.92
												±	±
Phyllostomidae	<i>G. soricina</i>	BZ	19	10.33 ± 1.87	Phytophagy	Gleaner	6.4	4.6	>1	Medium	10	99.41	117.86
												±	±
Phyllostomidae	<i>L. yerbabuena</i>	CMX	18	22.26 ± 1.72	Phytophagy	Gleaner	7	5.6	1	Large	15	24614.50	14780.52
												±	±
Phyllostomidae	<i>Lophostoma evotis</i>	BZ	1	19.00 ±	Animalivory	Gleaner	5.3	3	1	Small	20	3046.00	415.85
												±	±
Phyllostomidae	<i>M. waterhousii</i>	CMX	23	14.31 ± 2.63	Animalivory	Gleaner	5.8	5.6	1	Medium	10	616.37	415.85
												±	±
Phyllostomidae	<i>Mimon cozumelae</i>	BZ	3	25.00 ± 0	Animalivory	Gleaner	8.3	2.8	1	Small	20	2000.67	1284.91
												±	±
Phyllostomidae	<i>S. parvidens</i>	BZ	17	14.61 ± 1.55	Phytophagy	Gleaner	6.5	4	>1	Small	20	635.97	379.79
												±	±
Vespertilionidae	<i>Trachops cirrhosus</i>	BZ	3	31.50 ± 0.50	Animalivory	Gleaner	6.3	3.9	1	Medium	10	147.10	22.36
												±	±
Vespertilionidae	<i>A. pallidus</i>	MMX	10	15.80 ± 1.70	Animalivory	Gleaner	6.5	4	>1	Small	10	42.79	17.46
												±	±
Vespertilionidae	<i>E. furinalis</i>	BZ	7	7.57 ± 0.45	Animalivory	Aerial forager	6.2	3	>1	Large	20	36.66	40.54
												±	±
Vespertilionidae	<i>Lasiurus ega</i>	BZ	2	12.00 ± 4.24	Animalivory	Aerial forager	7.9	2.5	>1	Small	15	142.83	18.14
												±	±
Vespertilionidae	<i>M. velleri</i>	MMX	11	8.98 ± 1.02	Animalivory	Aerial forager	6.7	2.5	1	Large	10	211.65	170.79
												±	±

Capture sites: BZ, Belize; MMX, Northern Mexico; CMX, Central Mexico. N, number of hair sample analysed for cortisol. WAR, Wing Aspect Rati. Species were assigned to two dietary guilds: animalivory (insectivorous, sanguivorous and carnivorous) and phytophagy (frugivorous and nectarivorous species). Roost category assignment followed Patterson *et al.*, 2007.

(Dulude-de Broin *et al.*, 2019) but antibody dilution was adjusted. Antibody (R4866, C. Munro, University of California, Davis) and horseradish peroxidase dilutions were 1:10200 and 1:33400, respectively. Cortisol, rather than corticosterone, was targeted because it has been found to be the primary circulating GC in bats, with concentrations four times higher than corticosterone (reviewed in Kwiecinski and Damassa, 2000). Biochemical validation (parallelism and recovery) of the cortisol EIA was done using pooled hair extracts (see Supplementary data). We used pooled hair samples from Big Brown bats (*Eptesicus fuscus*) from a captive colony at McMaster University as a model for species with low cortisol concentrations (i.e. extracts reconstituted neat). For species with diluted extracts (*Tadarida brasiliensis*, *P. mexicanus* and *L. yerbabuena*) there was sufficient hair for separate pools. The inter-assay coefficient of variation (CV) for high control (24% binding) and low control (60% binding) were 9.3% and 9.8%, respectively. The intra-assay CV was 8.5%. The limits of detection and quantitation were 56 pg/ml and 153 pg/ml, respectively. Results are presented as nanograms of cortisol per gram of hair.

### Validation of immunoassay

Biochemical validations showed that the cortisol assay was suitable for hair. The recoveries of known concentrations of exogenous cortisol from hair extracts were  $87.1 \pm 3.5\%$  (dilution neat, *E. fuscus*),  $98.8 \pm 6.3\%$  (dilution, 1:5; *T. brasiliensis*),  $105.4 \pm 4.7\%$  (dilution, 1:5; *P. mexicanus*) and  $101.3 \pm 5.2\%$  (dilution, 1:50; *L. yerbabuena*). The measured hormone concentrations in the spiked samples correlated with the expected concentrations (*E. fuscus*:  $r = 0.997$ ,  $P < 0.001$ ; *T. brasiliensis*:  $r = 0.982$ ,  $P < 0.01$ ; *P. mexicanus*:  $r = 0.999$ ,  $P < 0.001$ ; *L. yerbabuena*:  $r = 0.995$ ,  $P < 0.001$ ; Supplementary data). Serial dilutions of pooled hair extracts showed parallel displacement with the cortisol standard curve (*E. fuscus*:  $r = 0.991$ ,  $P < 0.01$ ; *T. brasiliensis*:  $r = 0.988$ ,  $P < 0.01$ ; *P. mexicanus*:  $r = 0.997$ ,  $P < 0.001$ ; *L. yerbabuena*:  $r = 0.996$ ,  $P < 0.001$ ; Supplementary data).

### Species ecological traits

We compiled data on ecological traits considered relevant to cortisol mobilization from previously published literature and databases. Values for traits are species-level averages and may not reflect specific values at these sites (Table 1). Data on BMR was extracted from the literature (Cruz-Neto *et al.*, 2001; Genoud *et al.*, 2018; see Supplementary data) and when not available ( $n = 2$ ) the following formula was used for the estimation:  $\ln BMR = 0.744 \times \ln mass(\text{in } g) + 1.0895$  (Speakman and Thomas, 2003). Information on diet, foraging style, percentage of invertebrates in the diet and fecundity was extracted from the Elton Traits, PanTHERIA and Amniote Life History databases (Jones *et al.*, 2009b; Wilman *et al.*, 2014; Myhrvold *et al.*, 2015). We collapsed variation in diet into two dietary guilds: phytophagy (including nectarivores and frugivores) and animalivory (insectivores

and carnivores) because many bat species in our study have diets that combine more than one food source within these categories (Fenton *et al.*, 2001; Kunz and Fenton, 2005; Reid, 2009; Oelbaum *et al.*, 2019). We also considered the percentage of invertebrates in the diet of the animalivorous bats, which can vary significantly among species. Because foraging behaviour is a complex and plastic trait, we simplified this variable into two categories: aerial foragers (i.e. hawkers) and gleaners (including species that glean plant products like fruit as well as insects) since these behaviours may reflect differences in energetic demands associated with foraging (Herrera *et al.*, 2018). Because wing morphology can strongly influence the energetic costs of flight, we also included the mean wing aspect ratio for each species (Norberg and Rayner, 1987; Bullen *et al.*, 2014). Fecundity was defined as the annual average fecundity (litter size  $\times$  number of litters per year). We estimated roost durability following the methods of Patterson *et al.* (2007), where 1 indicates the most ephemeral and least protected roost types (e.g. rolled leaves and foliage) and 6 indicates the most permanent and protected roost types (e.g. caves). For species known to multiple use different kinds of roost, intermediate ranks were calculated, weighting roost categories according to the relative frequency of use reported in the literature (Schneeberger *et al.*, 2013). Lifespan was drawn from the Animal Ageing and Longevity database (AnAge: The Animal Ageing and Longevity Database, 2020) and DATLife (DATLife Database, 2020). Lifespan was grouped in five categories: 0–5, 5–10, 10–15, 15–20 and >20 years. For many of the species in these databases, longevity estimates are based on captive animals, which likely overestimates life expectancy in the wild. Because bats of a single species may live in colonies of varying sizes, and most values of colony size are reported in ranges in the literature, we classified maximum colony sizes reported for each species as small (1–50), medium (50–500) or large (>500) *sensu* Santana *et al.* (2011).

### Data analysis

We first used phylogenetic generalized least squares (PGLS) models to evaluate the effect of species-level ecological variables on hair cortisol concentrations while accounting for bat phylogenetic relatedness. We used the *rotl* and *ape* packages in R to extract the bat phylogeny from the Open Tree of Life and calculate branch lengths with Grafen's method (Paradis *et al.*, 2004; Michonneau *et al.*, 2016). We first fit a null PGLS model (intercept only) using the *nlme* package to estimate phylogenetic signal as Pagel's  $\lambda$  (Pagel, 1999). We next fit a PGLS model with bat family as the predictor to assess broad taxonomic patterns in hair cortisol. We then fit 15 PGLS univariate models with, dietary guild, foraging style, roost durability, fecundity, lifespan and colony size as predictors. We also fit five multivariate PGLS models including: BMR + body mass, dietary guild + fecundity, dietary guild + % invertebrates, dietary guild + lifespan + fecundity and dietary guild + fecundity + colony. We compared PGLS models with Akaike information criterion corrected for small


sample sizes (AICc) and assessed fit with an adjusted  $R^2$  (Burnham and Anderson, 2002). All PGLS models included weighting by sampling variance to account for variable sample sizes per species (Pennell, 2015).

We used generalized linear models (GLMs) to determine which individual- and habitat-level factors influence hair cortisol for each bat species. We first evaluated the relationship between body mass and hair cortisol separately for each species. Next, we ran species-specific GLMs including sex, reproductive stage (by sex) and site disturbance as predictors. Not all covariates were tested for all species due to sample size restrictions. Total sample size and balanced sample sizes among levels were considered to select the number of covariates to include in the model for each species. We included disturbance in GLMs only for species present in more than one site (*Pteronotus mesoamericanus*, *P. mexicanus*, *Macrotus waterhousii*, *T. brasiliensis*, *Glossophaga soricina*, *D. rotundus*) since disturbance was treated as constant within sites. The only genus sampled in both Belize and Mexico was *Pteronotus*. The two species *P. mesoamericanus* (Belize) and *P. mexicanus* (Mexico) represent lineages considered conspecific until a few years ago, but are now thought to represent distinct species that diverged very recently based on molecular and morphometric evidence (Pavan and Marroig, 2016). Because their phenotypes and ecology are still very similar, we treated these as conspecific to test if there were differences in hair cortisol between representatives from the two regions (Mexico and Belize). Tukey post hoc tests were conducted for significant covariates. We compared effect sizes across bat species by evaluating the degree of overlap in 95% confidence interval for each GLM coefficient. All analyses used the natural logarithm of hair cortisol as the response variable and assumed Gaussian errors. We confirmed that all models fulfilled assumptions of normality, homoscedasticity and non-multicollinearity (variance inflation factors < 3). We report data as mean  $\pm$  SD, unless otherwise noted.

## Results

### Ecological and evolutionary predictors of hair cortisol

We analysed 259 hair samples from 18 different bat species representing 5 families in Belize and Mexico (Table 1). Hair cortisol concentration across species varied by four orders of magnitude, ranging from  $36.6 \pm 40.5$  ng/g in *Eptesicus furinalis* to  $24\,614 \pm 14\,780$  ng/g in *L. yerbabuena* (Table 1). Even though mean hair cortisol apparently differed among families ( $F_{4,252} = 18.89$ ;  $P < 0.01$ ;  $R^2 = 0.23$ ; Fig. 2), this effect did not hold after accounting for phylogenetic relatedness ( $F_{4,13} = 1.84$ ;  $P = 0.18$ ;  $R^2 = 0.16$ ). Accordingly, we did not find strong phylogenetic signal in cortisol (Pagel  $\lambda = 0$ ). Ecological and life history traits were instead better predictors of species-level cortisol levels (Table 2). Mean hair cortisol was best predicted by a model including both dietary guild and fecundity; however, only fecundity had a significant effect



**Figure 2:** Mean and standard error of cortisol concentration in hair samples from 18 Neotropical bat species grouped by family. Summary statistics are displayed without adjusting for bat phylogeny. The y-axis is displayed on a  $\log_{10}$  scale.

( $F_{2,15} = 5.51$ ;  $P = 0.01$ ;  $R^2 = 0.34$ ; Table 2). Annual fecundity explained 24% of the variance in Neotropical bat mean hair cortisol. Species reported to have more than one pup per year had significantly lower cortisol than bats having only one pup per year ( $F_{1,16} = 6.22$ ;  $P = 0.02$ ; Fig. 3A). While phytophagous bats seem to have higher levels of cortisol in hair than animalivorous bats, this difference is not significant when considering the phylogenetic relatedness (Fig. 3B). Other ecological traits including roost durability, foraging style, and colony size were uninformative (Table 2). As cortisol levels varied by sex for some species (see below), we reran our model comparison after calculating species-level means for males and females separately. However, the above aggregate species-level results held when analysing the sexes separately (Supplementary Tables 2-3).


The lesser long-nosed bat (*L. yerbabuena*) showed particularly high hair cortisol ( $24\,614 \pm 14\,780$  ng/g). Because the high values of this species could bias inter-species comparisons, we assessed the sensitivity of our top models by excluding *L. yerbabuena*. In Fig. 3C, we show the coefficients from the top PGLS models with and without this species. In both cases, fecundity was the best species-level predictor of hair cortisol regardless of including *L. yerbabuena*.

### Individual-level analyses of bat hair cortisol

When investigating intra-specific variation, we found positive relationships between body mass and hair cortisol in two species: *P. mesoamericanus* ( $F_{1,24} = 7.34$ ;  $P = 0.010$ ;  $R^2 = 0.23$ ) and *Molossus nigricans* ( $F_{1,4} = 96.52$ ;  $P < 0.01$ ;  $R^2 = 0.96$ ; Fig. 4). The opposite trend was found in *P. mexicanus*, where heavier bats presented lower cortisol ( $F_{1,33} = 7.97$ ;  $P < 0.01$ ;  $R^2 = 0.19$ ). For *D. rotundus*, only sex was a significant

**Table 2:** PGLS models predicting hair cortisol (ln transformed) in Neotropical bats. Models are ranked by  $\Delta AICc$  with the number of coefficients ( $k$ ), Akaike weights ( $w_i$ ) and the adjusted  $R^2$

Model structure	df	$\Delta AICc$	$w_i$	$R^2$
Dietary guild + fecundity	2	0	0.412	0.34
Fecundity	4	1.08	0.247	0.24
Dietary guild+ fecundity+ foraging style	5	3.30	0.081	0.30
1 (intercept only)	6	4.12	0.054	0
Dietary guild + fecundity+ lifespan	1	4.38	0.048	0.46
Roost durability	2	5.64	0.025	0
Dietary guild	3	5.68	0.024	0.01
BMR+ body mass	3	6.00	0.021	0
Sample	2	6.39	0.017	0.33
Foraging style	2	6.67	0.015	0
Colony size	4	8.01	0.007	0.37
Dietary guild + invertebrate%	3	8.35	0.006	-0.03
BMR + foraging style	3	8.92	0.005	-0.08
Family	5	9.12	0.004	0.16
Foraging style + WAR	3	9.15	0.004	-0.1
Lifespan	4	10.54	0.002	-0.06




**Figure 3:** Left: Cortisol concentration in hair samples from 18 Neotropical bat species according to diet and annual fecundity. Y-axes are displayed on a  $\log_{10}$  scale. A. Mean hair cortisol by number of offspring per year. B. Hair cortisol by dietary guilds (animalivory or phytophagy). C. Differences in parameter estimates for the PGLS model with and without *L. yerbabuena*. Bars indicate the 95% confidence intervals. The reference values for each variable of the model are listed in parentheses.


predictor of hair cortisol ( $F_{1,19} = 4.39$ ;  $P = 0.04$ ;  $R^2 = 0.19$ ). Male vampire bats had significantly lower hair cortisol than females ( $t_{20} = 2.09$ ;  $P = 0.02$ ). Similarly, variation in hair cortisol in the moustached bat (*P. mesoamericanus*) and the mastiff bat (*M. nigricans*) was explained only by sex, with males having lower concentrations than females ( $t_4 = 2.68$ ;  $P = 0.01$  and  $t_{24} = -6.373$ ;  $P = 0.01$ , respectively; Fig. 5). When treating *P. mesoamericanus* (Belize) and *P. mexicanus*

(Mexico) as one species, we found differences in hair cortisol between the two populations. Bats from Mexico had higher cortisol than their counterparts in Belize ( $F_{1,59} = 29.88$ ;  $P < 0.01$ ;  $R^2 = 0.33$ ). Within Mexico, hair cortisol in *P. mexicanus* was explained by site disturbance ( $F_{2,31} = 72.35$ ;  $P < 0.001$ ): bats roosting in Don Pancho cave (San Agustin island), a site with moderate disturbance (HMI = 0.38), showed significantly higher hair cortisol than bats roosting in





**Figure 4:** Relationship between hair cortisol concentration and body mass for each Neotropical bat species. Lines represent the GLM fit for each species. The y-axis is shown on a log<sub>10</sub> scale.



**Figure 5:** Mean and standard error for hair cortisol concentration by sex (F, female; M, male) for nine species of Neotropical bat species. Asterisks indicate species for which cortisol significantly varies by sex. Effects are only shown for species with balanced sample sizes per sex. Y-axes are shown on a log<sub>10</sub> scale.

## Discussion

Analysis of hair cortisol has become a popular method to study long-term stress in wild animals, offering several practical advantages (e.g. minimally invasive collection, easy sample storage and transport). An accurate interpretation of cortisol levels attributed to stress, however, requires a good understanding of the intrinsic and extrinsic drivers of baseline variation. Factors influencing hair cortisol in bats must be identified before hair cortisol can be used as a conservation tool to assess effects of environmental conditions on bat population health. In this study, we present the first quantification of hair cortisol in bats and describe relationships between hair cortisol levels and both intrinsic and ecological traits. Cortisol in blood, faeces and hair are known to be highly correlated in various mammals (e.g. chimpanzees, chipmunks and mice; Kalliokoski *et al.*, 2019). Therefore, although the concentration values are not directly comparable across different matrices, the effects of covariates can still be compared with our results.

One of the advantages of the use of hair as a physiological biomarker of stress is the longer window of time that it provides compared to other samples like blood and faeces. This feature, however, is only useful if the rate of hair growth and moulting patterns are known for the species of concern. Moulting cycles in bats are understudied, especially in tropical bats (Fraser *et al.*, 2013). Therefore, we cannot be certain of the precise temporal window that this biomarker could represent for our study species. While moulting patterns for most of our tropical study species are unknown, we know that many temperate species have a one annual summer/fall moult cycle (Fraser *et al.*, 2013). If we assume our study species have a similar cycle, then the hair cortisol levels detected in our study would be reflecting primarily the circulating cortisol levels during one to two months of rapid hair growth between July and September. Moulting in tropical bats, however,

El Salitre and La Fabrica caves in Colima ( $t_{20} = 9.94, P < 0.01$ ;  $t_{21} = -10.29$ ; Fig. 1). There was no effect of sex ( $t_{33} = -1.15$ ;  $P > 0.31$ ) or females' reproductive stage ( $F_{4,14} = 3.26$ ;  $P = 0.06$ ) on hair cortisol in *P. mexicanus*. For other species such as *E. furinalis* ( $F_{1,12} = 2.451$ ;  $P = 0.64$ ), *L. yerbabuena* ( $F_{1,17} = 0.52$ ;  $P = 0.94$ ), *Saccopteryx billineata* ( $F_{2,17} = 0.18$ ;  $P = 0.83$ ), *Rhynchonycteris naso* ( $F_{2,11} = 2.60$ ;  $P = 0.12$ ), *G. soricina* ( $F_{3,15} = 0.13$ ;  $P = 0.94$ ), *Macrotus waterhousii* ( $F_{2,18} = 0.24$ ;  $P = 0.78$ ), *Sturnira parvidens* ( $F_{2,14} = 0.2052$ ,  $P = 0.81$ ) and *A. pallidus* ( $F_{2,9} = 0.506$ ;  $P = 0.68$ ), none of the individual- or habitat-level traits examined were informative predictors of hair cortisol levels.

might not be confined to a distinct or single annual event, given the relative stable environmental conditions the tropics offer. Moulting patterns in tropical bats therefore should be investigated further to allow inferences about the window of time over which cortisol levels in hair integrate and reflect circulating levels. Other aspects associated with differences in the timing of new fur growth, and therefore cortisol deposition in hair, could be related to local environmental variables (e.g. precipitation, temperature and food seasonality; *Tiunov and Makarikova, 2007*), as well as species-specific life history traits (e.g. reproduction and migration; *Heydon et al., 1995*).

Overall, we found particularly high hair cortisol in most bat species compared to levels reported in hair for other small mammals (e.g. chipmunk [66–110 g] = 40.27–260.22 ng/g of hair; *Mastromonaco et al., 2014*). Previous studies in bats, examining plasma and faeces, have also reported higher cortisol levels relative to similar samples from other mammal species (*Lewanzik et al., 2012*; *Kelm et al., 2016*; *Hald, 2019*). Some of the exceptional life history traits of bats, such as long lifespan and low fecundity, could explain why bats exhibit higher levels of GCs compared to other mammals (*Austad and Fischer, 1991*). According to life history theory, long-lived species with low reproduction rates are expected to prioritize their adult survival (i.e. future offspring) over current reproduction (*Stearns, 1992*), which could in turn favour higher investment in self-maintenance that might be facilitated by high baseline levels of GCs (*Ricklefs and Wikelski, 2002*).

### Ecological factors among species

Despite the similarity in the HPA hormonal pathways across vertebrates, baseline and stress-induced GC levels are context and species specific (*Romero, 2004*; *French et al., 2008*; *Crespi et al., 2013*; *Kalliokoski et al., 2019*). In light of this, it is not surprising that hair cortisol levels in bats showed broad interspecific variation. The differences found could not be explained solely by taxonomic family or phylogenetic relatedness ( $\lambda = 0$ ), which suggests that other environmental and ecological factors are influencing hair cortisol in Neotropical bats.

Among all the ecological traits evaluated, annual fecundity was the best predictor of hair cortisol. Species with lower fecundity showed higher concentrations of cortisol in hair. This relationship can be supported from a physiological perspective, considering that GCs play an important role modulating the production of reproductive hormones upstream in the hypothalamic–pituitary–gonadal axis, both under homeostatic and challenging conditions. This interaction, however, has only been clearly demonstrated within populations (e.g. individuals with high levels of cortisol reduce reproduction, reallocating energy current needs). It is unknown if interspecific variation in GCs is driven by the same mechanisms. Looking to other taxa, studies on birds have found similar results to those here in Neotropical bats, where avian species with low clutch size and few breeding events showed higher circulating GCs (*Bókony et al., 2009*; *Ouyang et al., 2011*).

This pattern is supported by predictions from life history theory; for species with lower fecundity, the value of each offspring is higher than in species with relatively high fecundity (*Lendvai et al., 2007*). Parents of more valuable broods would be predicted to be more ‘willing’ to invest in offspring survival, which might be facilitated by high baseline GC levels (*Bókony et al., 2009*). Confirming if this mechanism explains the GCs levels found in our bat species would require more information about reproductive strategies (e.g. monoestry and polyestry) and seasonality (*Wingfield and Sapolsky, 2003*), which is limited for most Neotropical species. A more practical rationale for our results could be related to differences in hair cortisol deposition among species with different reproductive cycles. For many species, pelage moulting occurs after breeding (*Constantine, 1957*; *Dwyer, 1963*; *Cryan et al., 2004*; *Measor et al., 2017*), given that both reproduction and hair growth are energetically demanding processes (*Ling, 1970*). Based on this pattern, we could expect that cortisol deposition in hair may reflect differences in breeding events among bat species.

Although diet explained additional variation in Neotropical bat hair cortisol, this variable was uninformative when considering phylogenetic relationships; hair cortisol did not vary significantly between our simplified dietary guilds. Further studies using a more accurate classification of diet (e.g. using stable isotopes or metabarcoding; *Oelbaum et al., 2019*; *Ingala et al., 2021*) could give more conclusive insights into the links between feeding strategies and cortisol levels in bats.

GCs play a key role in metabolic function, facilitating fuel mobilization (e.g. glucose, fatty acids) under normal and challenging conditions (*Kuo et al., 2015*). A positive relationship between resting metabolic rate (RMR) and plasma cortisol levels has been reported for various mammalian species (including four species of bats), and this relationship has been suggested as a general pattern for mammals (*Haase et al., 2016*). Due to the limited data on RMR for our study species, we used BMR as an indicator of energy expenditure. Different to what we expected, BMR was not an informative factor for cortisol variation in hair among the bats in our study. The positive relationship between cortisol levels and metabolic rate previously found in plasma might be obscured in studies of hair cortisol like ours, due to confounding factors such as moulting cycles and cortisol deposition rate. In addition, obtaining accurate BMRs in wildlife species (particularly free-ranging animals) is challenging, which raises questions about the quality of BMR data, especially in comparative studies (*Genoud et al., 2018*). For future studies, a more realistic and informative indicator of energy turnover in free-ranging animals is the Daily Energy Expenditure (*Speakman, 1997*), which integrates the energy allocated in different activities such as foraging, commuting and thermoregulation (*Butler et al., 2004*).

The relationship between body condition and GC release has been widely evaluated, because weight loss is one of the early responses to long-term stress in many species (*Kitaysky*

*et al.*, 1999; Angelier *et al.*, 2009; Dickens and Romero, 2013). However, the direction of the effect of body condition on cortisol is context and species dependent (Crespi *et al.*, 2013). We used body mass as an indicator of body condition because it is a more informative metric than other indices in bats (McGuire *et al.*, 2018). We found different directions of the effect of body mass on hair cortisol. For two of the studied species (*M. nigricans* and *P. mesoamericanus*), heavier individuals showed higher concentrations of hair cortisol. In contrast, *P. mexicanus* showed a negative relationship between body mass and cortisol. These divergent results suggest that the relationship between cortisol and body condition of bats is not generally predictable and might be species specific. Hair cortisol only reflects the time window of fur growth. Therefore, it is possible that body condition might have changed since the individual's last moult.

One species that stood out for its particularly high levels of cortisol was *L. yerbabuena*. This species is highly mobile and migratory (Horner *et al.*, 1998; Buecher and Sidner, 2013; Medellín *et al.*, 2018). Migration itself was not considered in our analyses, because the degree to which bats may migrate seasonally is unclear for many of the species in our sample. Migratory behaviour, however, could explain such high cortisol concentrations in *L. yerbabuena*. La Fábrica caves in Colima, one of our field sites, is known to be one of the starting points of the annual migration of *L. yerbabuena* (Medellín *et al.*, 2018). The role of GCs during migration has been widely studied in birds, fish and some large mammals, but not in bats (Holberton, 1999; Romero, 2002; Wada, 2008). We hypothesize that premigratory fattening could explain the high hair cortisol levels observed in *L. yerbabuena*, and we encourage future studies to address this question.

Consistent with other studies, we found differences in hair cortisol levels between sexes, albeit for only 4 of our 18 studied species: *D. rotundus*, *M. nigricans*, *P. mexicanus* and *P. mesoamericanus*. For these species, females showed higher cortisol than males, a trend that appears to hold for many mammalian species (Bechshøft *et al.*, 2011; Hau *et al.*, 2016; Rakotoniaina *et al.*, 2017; Dettmer *et al.*, 2018). Higher levels of GCs in females can be attributed to sex differences in the HPA axis activity, which are mainly mediated by gonadal steroid hormones (i.e. androgens and estrogens). For example, estradiol, which is more abundant in females than males, can enhance cortisol release, while androgens tend to reduce its production (Handa *et al.*, 1994). Other mechanisms underlying sex differences in HPA axis regulation and stress-related behaviours in mammals are reviewed by Zuloaga *et al.* (2020). Females have also shown differences in HPA axis activity depending on their life history stage, with GCs being higher during the late stages of pregnancy (Reeder *et al.*, 2004). Studies in a fruit-eating bat (*Artibeus jamaicensis*) and little brown myotis (*Myotis lucifugus*) have reported higher levels of plasma GCs in pregnant females (Reeder and Kramer, 2005; Klose *et al.*, 2006). Contrary to those findings, we did not find reproductive state to influence hair cortisol in

our female-only model (i.e. for *P. mesoamericanus* in Belize). However, it may have been difficult to detect an effect, given the low number of pregnant females in our sample ( $n = 5$ , 24%) and the fact that moulting might not occur in conjunction with mating. Although cortisol has been proposed as the primary GC in bats (reviewed by Kwiecinski and Damassa, 2000), corticosterone is also detectable in circulation and has been identified to play an important role in reproduction (Koren *et al.*, 2012). Therefore, some effects of ecological traits could go unnoticed and the complexity of the stress response in bats could be oversimplified by quantifying only cortisol.

Bats have been proposed as good indicators of habitat quality due to their ecological diversity, wide distribution and potential sensitivity to disturbance (Jones *et al.*, 2009a; Cunto and Bernard, 2012; Stahlschmidt and Brühl, 2012). However, a clear correlation between environmental disturbance and cortisol levels, in faeces and blood, has not been reported in bats (Wada *et al.*, 2010; Allen *et al.*, 2011; Kelm *et al.*, 2016). Cortisol in hair could reflect better the effects of chronic stressors such as human settlements, and it is not sensitive to capture stress. We compared hair cortisol in three of our study species found in sites with varying fragmentation and agricultural activities in Mexico. We found an effect of disturbance in only one of these species, *P. mexicanus*, for which bats roosting in Don Pancho Cave island, a site with moderate disturbance, showed the highest concentrations of cortisol (Fig. 1). We speculate that the high levels found in this population could reflect differences in the cave microhabitat compared to the other caves in our Mexican sample. Don Pancho Cave is a narrow crevice estimated to have a higher colony size (100 000 individuals from 6 species; Téllez *et al.*, 2018) than the other sampled caves El Salitre (~10 000 individuals from 10 species; Torres-Flores *et al.*, 2012) and La Fábrica (>5000 individuals from four species). The high density of bats in the Don Pancho cave may increase agonistic social interactions (Creel *et al.*, 2013) and parasite transmission (Langwig *et al.*, 2012; Postawa and Szubert-Kruszyńska, 2014), factors that have been shown to increase cortisol levels in other mammals.

Physiological responses to chronic stress in wildlife are difficult to unravel and predict unless multiple responses at different levels of biological organization are evaluated simultaneously (Dickens and Romero, 2013). Hair cortisol offers great potential as a tool to monitor health in wild populations, particularly those already identified at risk (Kalliokoski *et al.*, 2019). For instance, chronically elevated cortisol levels have been linked to greater susceptibility to infection and disease severity (Davy *et al.*, 2017). Periodic surveys of hair cortisol could therefore help identify periods when bats might be more vulnerable to infection (e.g. white nose syndrome). Further, such surveys might also inform when individuals are more likely to shed zoonotic pathogens (e.g. henipaviruses and filoviruses; Plowright *et al.*, 2008; Davy *et al.*, 2017; McMichael *et al.*, 2017; Kessler *et al.*, 2018).

## Conclusions

The current study reports cortisol levels in hair of 18 Neotropical bat species from two countries and serves as a reference for future research using this method in wild bat populations. We found that fecundity and potentially diet are important ecological traits explaining interspecific variation in bat hair cortisol. Within species, female bats exhibited higher cortisol than males and the effect of body mass varied among species. Other factors that may be important at the individual level, such as parasite load and colony size, should be considered in future studies to have a more complete understanding of sources of variation on baseline GC levels within species. Importantly, studies looking at hair growth rate and moulting cycles in Neotropical bat species are imperative to give an accurate interpretation of hair cortisol as a biomarker of stress response. Applied properly, hair cortisol quantification is a powerful minimally invasive technique with multiple potential applications in bat ecology, physiology and conservation. Our findings and ongoing work will help to validate and apply hair cortisol as a monitoring tool in wild bat populations.

## Funding

This work was supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant (to K.C.W.), the Toronto Zoo Foundation (to G.M.), the ARCS Foundation and the American Museum of Natural History Theodore Roosevelt Memorial Fund (to D.J.B.) and the Taxonomic Mammalogy Fund of the American Museum of Natural History (to N.B.S.).

## Author contributions statement:

N.S.H. conceived the study and collected the samples. G.F.M. analysed the samples. N.S.H. and D.J.B. analysed the data and wrote the manuscript. N.B.S. provided support for collection and export permits and coordinated field work logistics in Belize. K.C.W. Jr. secured funding for field and laboratory work. All authors reviewed drafts of the manuscript.

## Acknowledgments

We thank Christine Gilman, Patricia Medd and Paula Mackie for their assistance with the cortisol assays. We also thank Brock Fenton, Sara Ketelsen, Neil Duncan, the numerous members of the Lamanai bat research team and the staff of the Lamanai Field Research Center for their assistance with bat capture, field logistics and permits in Belize.

## Supplementary material

Supplementary material is available at *Conservation Physiology* online.

## References

- Acker M, Mastromonaco G, Schulte-Hostedde AI (2018) The effects of body region, season and external arsenic application on hair cortisol concentration. *Conserv Physiol* 6: coy037.
- Allen LC, Turmelle AS, Widmaier EP, Hristov NI, McCracken GF, Kunz TH (2011) Variation in physiological stress between bridge- and cave-roosting Brazilian free-tailed bats. *Conserv Biol* 25: 374–381.
- AnAge: The Animal Ageing and Longevity Database (2020). <https://genomics.senescence.info/species/> (last accessed 4 March 2020).
- Andreasson U, Perret-Liaudet A, van Waalwijk van Doorn LJC, Blennow K, Chiasserini D, Engelborghs S, Fladby T, Genc S, Kruse N, Kuiperij HB, et al. (2015) A practical guide to immunoassay method validation. *Front Neurol* 6: 179.
- Angelier F, Holberton RL, Marra PP (2009) Does stress response predict return rate in a migratory bird species? A study of American redstarts and their non-breeding habitat. *Proc R Soc B Biol Sci* 276: 3545–3551.
- Ashley NT, Barboza PS, Macbeth BJ, Janz DM, Cattet MRL, Booth RK, Wasser SK (2011) Glucocorticosteroid concentrations in feces and hair of captive caribou and reindeer following adrenocorticotrophic hormone challenge. *Gen Comp Endocrinol* 172: 382–391.
- Austad SN, Fischer KE (1991) Mammalian aging, metabolism, and ecology: evidence from the bats and marsupials. *J Gerontol* 46: 47–53.
- Azevedo A, Bailey L, Bandeira V, Dehnhard M, Fonseca C, De Sousa L, Jewgenow K (2019) Age, sex and storage time influence hair cortisol levels in a wild mammal population. *PLoS One* 14: 8. [10.1371/journal.pone.0221124](https://doi.org/10.1371/journal.pone.0221124).
- Azevedo A, Wauters J, Kirschbaum C, Serra R, Rivas A, Jewgenow K (2020) Sex steroids and glucocorticoid ratios in Iberian lynx hair. *Conserv Physiol* 8: 75.
- Bechshøft T, Sonne C, Dietz R, Born EW, Novak MA, Henchey E, Meyer JS (2011) Cortisol levels in hair of East Greenland polar bears. *Sci Total Environ* 409: 831–834.
- Becker DJ, Chumchal MM, Broders HG, Korstian JM, Clare EL, Rainwater TR, Platt SG, Simmons NB, Fenton MB (2018) Mercury bioaccumulation in bats reflects dietary connectivity to aquatic food webs. *Environ Pollut* 233: 1076–1085.
- Bókony V, Lendvai ÁZ, Likér A, Angelier F, Wingfield JC, Chastel O (2009) Stress response and the value of reproduction: are birds prudent parents? *Am Nat* 173: 589–598.
- Bonier F, Martin PR (2016) How can we estimate natural selection on endocrine traits? Lessons from evolutionary biology. *Proc R Soc B Biol Sci* 283. [10.1098/rspb.2016.1887](https://doi.org/10.1098/rspb.2016.1887).

- Bonier F, Martin PR, Moore IT, Wingfield JC (2009) Do baseline glucocorticoids predict fitness? *Trends Ecol Evol* 24: 634–642.
- Boonstra R (2013) The ecology of stress: a marriage of disciplines. *Funct Ecol* 27: 7–10.
- Buecher DC, Sidner R (2013) Long distance commutes by lesser long-nosed bats (*Leptonycteris yerbabuena*) to visit residential hummingbird feeders. In *Merging Science and Management in a Rapidly Changing World: Biodiversity and Management of the Madrean Archipelago III and 7th Conference on Research and Resource Management in the Southwestern Deserts*, 1–5 May 2012. Tucson, AZ. Proceedings. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO, pp. 427–433.
- Bullen RD, McKenzie NL, Cruz-Neto AP (2014) Aerodynamic power and mechanical efficiency of bat airframes using a quasi-steady model. *CEAS Aeronaut J* 5: 253–264.
- Burnham K, Anderson D (2002) A practical information-theoretic approach. In *Model Selection and Multimodel Inference*, Ed 2. Springer, New York.
- Busch DS, Hayward LS (2009) Stress in a conservation context: a discussion of glucocorticoid actions and how levels change with conservation-relevant variables. *Biol Conserv* 142: 2844–2853.
- Butler PJ, Green JA, Boyd IL, Speakman JR (2004) Measuring metabolic rate in the field: the pros and cons of the doubly labelled water and heart rate methods. *Funct Ecol* 18: 168–183.
- Constantine DG (1957) Color variation and molt in *Tadarida brasiliensis* and *Myotis velifer*. *J Mammal* 38: 461.
- Creagh BW, Brendan Delehanty RB (2013) Reality as the leading cause of stress: rethinking the impact of chronic stress in nature. *Funct Ecol* 27: 11–23.
- Creel S, Dantzer B, Goymann W, Rubenstein DR (2013) The ecology of stress: effects of the social environment. *Funct Ecol* 27: 66–80.
- Crespi EJ, Williams TD, Jessop TS, Delehanty B (2013) Life history and the ecology of stress: how do glucocorticoid hormones influence life-history variation in animals? *Funct Ecol* 27: 93–106.
- Cruz-Neto AP, Garland T, Abe AS (2001) Diet, phylogeny, and basal metabolic rate in phyllostomid bats. *Fortschr Zool* 104: 49–58.
- Cryan PM, Bogan MA, Rye RO, Landis GP, Kester CL (2004) Stable hydrogen isotope analysis of bat hair as evidence for seasonal molt and long-distance migration. *J Mammal* 85: 995–1001.
- Cunto GC, Bernard E (2012) Neotropical bats as indicators of environmental disturbance: what is the emerging message? *Acta Chiropt* 14: 143–151.
- DATLife Database. Max-Planck Institute for Demographic Research (Germany) (2020). <https://datlife.org/> (last accessed 4 March 2020).
- Davenport MD, Tiefenbacher S, Lutz CK, Novak MA, Meyer JS (2006) Analysis of endogenous cortisol concentrations in the hair of rhesus macaques. *Gen Comp Endocrinol* 147: 255–261.
- Davy CM, Mastro Monaco GF, Riley JL, Baxter-Gilbert JH, Mayberry H, Willis CKR (2017) Conservation implications of physiological carry-over effects in bats recovering from white-nose syndrome. *Conserv Biol* 31: 615–624.
- Dettmer AM, Murphy AM, Guitarra D, Slonecker E, Suomi SJ, Rosenberg KL, Novak MA, Meyer JS, Hinde K (2018) Cortisol in neonatal mother's milk predicts later infant social and cognitive functioning in rhesus monkeys. *Child Dev* 89: 525–538.
- Dettmer AM, Novak MA, Meyer JS, Suomi SJ (2014) Population density-dependent hair cortisol concentrations in rhesus monkeys (*Macaca mulatta*). *Psychoneuroendocrinology* 42: 59–67.
- Dickens MJ, Romero LM (2013) A consensus endocrine profile for chronically stressed wild animals does not exist. *Gen Comp Endocrinol* 191: 177–189.
- Dulude-de Broin F, Côté SD, Whiteside DP, Mastro Monaco GF (2019) Faecal metabolites and hair cortisol as biological markers of HPA-axis activity in the rocky mountain goat. *Gen Comp Endocrinol* 280: 147–157.
- Dumont ER, Dávalos LM, Goldberg A, Santana SE, Rex K, Voigt CC (2012) Morphological innovation, diversification and invasion of a new adaptive zone. *Proc R Soc B Biol Sci* 279: 1797–1805.
- Dwyer PD (1963) Seasonal changes in pelage of *Miniopterus schreibersi blepotis* (Chiroptera) in north-eastern New South Wales. *Aust J Zool* 11: 290–300.
- Fenton MB (1990) The foraging behaviour and ecology of animal-eating bats. *Can J Zool* 68: 411–422.
- Fenton MB, Bernard E, Bouchard S, Hollis L, Johnston DS, Lausen CL, Ratcliffe JM, Riskin DK, Taylor JR, Zigouris J (2001) The bat fauna of Lamanai, Belize: roosts and trophic roles. *J Trop Ecol* 17: 511–524.
- Fourie NH, Brown JL, Jolly CJ, Phillips-Conroy JE, Rogers J, Bernstein RM (2016) Sources of variation in hair cortisol in wild and captive non-human primates. *Fortschr Zool* 119: 119–125.
- Fraser EE, Longstaffe FJ, Fenton MB (2013) Moulting matters: the importance of understanding moulting cycles in bats when using fur for endogenous marker analysis. *Can J Zool* 91: 533–544.
- Fraser KC, McKinnon EA, Diamond AW (2010) Migration, diet, or molt? interpreting stable-hydrogen isotope values in neotropical bats. *Biotropica* 42: 512–517.
- French SS, Fokidis HB, Moore MC (2008) Variation in stress and innate immunity in the tree lizard (*Urosaurus ornatus*) across an urban-rural gradient. *J Comp Physiol B Biochem Syst Environ Physiol* 178: 997–1005.
- Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, Turner GG, Butchkoski CM, Kunz TH (2010) An emerging disease causes regional population collapse of a common North American bat species. *Science* 329: 679–682.
- Genoud M, Isler K, Martin RD (2018) Comparative analyses of basal rate of metabolism in mammals: data selection does matter: comparative analyses of basal rate of metabolism. *Biol Rev* 93: 404–438.

- Gunnell GF, Simmons NB (2012) *Evolutionary History of Bats. Evolutionary History of Bats: Fossils, Molecules and Morphology*. Cambridge University Press, Cambridge.
- Haase CG, Long AK, Gillooly JF (2016) Energetics of stress: linking plasma cortisol levels to metabolic rate in mammals. *Biol Lett* 12: 20150867. [10.1098/rsbl.2015.0867](https://doi.org/10.1098/rsbl.2015.0867).
- Hald HA (2019) *Cortisol Measurements in Female Whiskered Bats (Myotis mystacinus) During an Extraordinarily Warm and Dry Norwegian Summer*. (Master's dissertation). Norwegian University of Science and Technology, Trondheim, Norway.
- Handa RJ, Nunley KM, Lorens SA, Louie JP, McGivern RF, Bollnow MR (1994) Androgen regulation of adrenocorticotropic and corticosterone secretion in the male rat following novelty and foot shock stressors. *Physiol Behav* 55: 117–124.
- Hau M, Casagrande S, Ouyang JQ, Baugh AT (2016) Glucocorticoid-mediated phenotypes in vertebrates: multilevel variation and evolution. *Adv Study Behav* 48: 41–115.
- Hernout BV, McClean CJ, Arnold KE, Walls M, Baxter M, Boxall ABA (2016) Fur: a non-invasive approach to monitor metal exposure in bats. *Chemosphere* 147: 376–381.
- Herrera JP, Duncan N, Clare E, Fenton MB, Simmons N (2018) Disassembly of fragmented bat communities in Orange Walk District, Belize. *Acta Chiroptol* 20: 147.
- Heydon MJ, Milne JA, Brinklow BR, Loudon ASI (1995) Manipulating melatonin in red deer (*Cervus elaphus*): differences in the response to food restriction and lactation on the timing of the breeding season and prolactin-dependent pelage changes. *J Exp Zool* 273: 12–20.
- Hing S, Narayan EJ, Thompson RCA, Godfrey SS (2016) The relationship between physiological stress and wildlife disease: consequences for health and conservation. *Wildl Res* 43: 51–60.
- Holberton RL (1999) Changes in patterns of corticosterone secretion concurrent with migratory fattening in a neotropical migratory bird. *Gen Comp Endocrinol* 116: 49–58.
- Horner MA, Fleming TH, Sahey CT (1998) Foraging behaviour and energetics of a nectar-feeding bat, *Leptonycteris curasoae* (Chiroptera: Phyllostomidae). *J Zool* 244: 575–586.
- INEGI (2013) Uso de suelo y vegetación. *Datos vectoriales escala 1250 000 Ser V* 1–84.
- Ingala MR, Simmons NB, Wulsch C, Krampis K, Provost KL, Perkins SL (2021) Molecular diet analysis of neotropical bats based on fecal DNA metabarcoding. *Ecol Evol* 00: ece3.7579.
- IUCN *Red List of Threatened Species* (2020). <https://www.iucnredlist.org/search?query=bats&searchType=species> (last accessed 4 July 2020).
- Jones G, Jacobs DS, Kunz TH, Wilig MR, Racey PA (2009a) Carpe noctem: the importance of bats as bioindicators. *Endanger Species Res* 8: 93–115.
- Jones KE, Bielby J, Cardillo M, Fritz SA, O'Dell J, Orme CDL, Safi K, Sechrest W, Boakes EH, Carbone C *et al.* (2009b) PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* 90: 2648–2648.
- Kalliokoski O, Jellestad FK, Murison R (2019) A systematic review of studies utilizing hair glucocorticoids as a measure of stress suggests the marker is more appropriate for quantifying short-term stressors. *Sci Rep* 9: 1–14.
- Kelm DH, Popa-Lisseanu AG, Dehnhard M, Ibáñez C (2016) Non-invasive monitoring of stress hormones in the bat *Eptesicus isabellinus*—do fecal glucocorticoid metabolite concentrations correlate with survival? *Gen Comp Endocrinol* 226: 27–35.
- Kennedy CM, Oakleaf JR, Theobald DM, Baruch-Mordo S, Kiesecker J (2019) Managing the middle: a shift in conservation priorities based on the global human modification gradient. *Glob Chang Biol* 25: 811–826.
- Kessler MK, Becker DJ, Peel AJ, Justice NV, Lunn T, Crowley DE, Jones DN, Eby P, Sánchez CA, Plowright RK (2018) Changing resource landscapes and spillover of henipaviruses. *Ann N Y Acad Sci* 1429: 78–99.
- Kitaysky AS, Wingfield JC, Piatt JF (1999) Dynamics of food availability, body condition and physiological stress response in breeding black-legged Kittiwakes. *Funct Ecol* 13: 577–584.
- Klose SM, Smith CL, Denzel AJ, Kalko EKV (2006) Reproduction elevates the corticosterone stress response in common fruit bats. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 192: 341–350.
- Koren L, Mokady O, Karaskov T, Klein J, Koren G, Geffen E (2002) A novel method using hair for determining hormonal levels in wildlife. *Anim Behav* 403–406.
- Koren L, Whiteside D, Fahlman Å, Ruckstuhl K, Kutz S, Checkley S, Dumond M, Wynne-Edwards K (2012) Cortisol and corticosterone independence in cortisol-dominant wildlife. *Gen Comp Endocrinol* 177: 113–119.
- Książek A, Zub K, Szafrńska PA, Wieczorek M, Konarzewski M (2017) The nexus of hair corticosterone level, immunocompetence, metabolic rates and overwinter survival in the root vole, *Microtus oeconomus*. *Gen Comp Endocrinol* 250: 46–53.
- Kunz T, Fenton M (2005) *Bat Ecology*. University of Chicago Press, Chicago, USA.
- Kunz TH, Arnett EB, Erickson WP, Hoar AR, Johnson GD, Larkin RP, Strickland MD, Thresher RW, Turtle MD (2007) Ecological impacts of wind energy development on bats: questions, research needs, and hypotheses. *Front Ecol Environ* 5: 315–324.
- Kunz TH, Parsons S (2009) *Ecological and Behavioral Methods for the Study of Bats*. Johns Hopkins University Press.
- Kuo T, McQueen A, Chen TC, Wang JC (2015) Regulation of glucose homeostasis by glucocorticoids. *Adv Exp Med Biol* 872: 99–126.
- Kwiecinski G, Damassa D (2000) Peripheral endocrines in bat reproduction. In E Crichton, P Krutzsch, eds, *Reproductive Biology of Bats*. Academic Press, New York, NY, pp. 65–89.

- Landys MM, Ramenofsky M, Wingfield JC (2006) Actions of glucocorticoids at a seasonal baseline as compared to stress-related levels in the regulation of periodic life processes. *Gen Comp Endocrinol* 148: 132–149.
- Langwig KE, Frick WF, Bried JT, Hicks AC, Kunz TH, Marm Kilpatrick A (2012) Sociality, density-dependence and microclimates determine the persistence of populations suffering from a novel fungal disease, white-nose syndrome. *Ecol Lett* 15: 1050–1057.
- Lavergne SG, Peers MJL, Mastromonaco G, Majchrzak YN, Nair A, Boutin S, Boonstra R (2020) Hair cortisol as a reliable indicator of stress physiology in the snowshoe hare: influence of body region, sex, season, and predator–prey population dynamics. *Gen Comp Endocrinol* 294. [10.1016/j.ygcen.2020.113471](https://doi.org/10.1016/j.ygcen.2020.113471).
- Lendvai ÁZ, Giraudeau M, Chastel O (2007) Reproduction and modulation of the stress response: an experimental test in the house sparrow. *Proc R Soc B Biol Sci* 274: 391–397.
- Lewanzik D, Kelm DH, Greiner S, Dehnhard M, Voigt CC (2012) Ecological correlates of cortisol levels in two bat species with contrasting feeding habits. *Gen Comp Endocrinol* 177: 104–112.
- Ling JK (1970) Pelage and molting in wild mammals with special reference to aquatic forms. *Q Rev Biol* 45: 16–54.
- Macbeth BJ, Cattet MRL, Obbard ME, Middel K, Janz DM (2012) Evaluation of hair cortisol concentration as a biomarker of long-term stress in free-ranging polar bears. *Wildl Soc Bull* 36: 747–758.
- Macbeth BJ, Cattet MRL, Stenhouse GB, Gibeau ML, Janz DM (2010) Hair cortisol concentration as a noninvasive measure of long-term stress in free-ranging grizzly bears (*Ursus arctos*): considerations with implications for other wildlife. *Can J Zool* 88: 935–949.
- MacDougall-Shackleton SA, Bonier F, Romero LM, Moore IT (2019) Glucocorticoids and “stress” are not synonymous. *Integr Org Biol* 1. [10.1093/iob/obz017](https://doi.org/10.1093/iob/obz017).
- Magioli M, Bovo AAA, Alberici V, De Barros Ferraz KMPM (2019) The use of hair traps as a complementary method in mammal ecology studies. *Mammalia* 83: 144–149.
- Mastromonaco GF, Gunn K, McCurdy-Adams H, Edwards DB, Schulte-Hostedde AI (2014) Validation and use of hair cortisol as a measure of chronic stress in eastern chipmunks (*Tamias striatus*). *Conserv Physiol* 2. [10.1093/conphys/cou055](https://doi.org/10.1093/conphys/cou055).
- McGuire LP, Kelly LA, Baloun DE, Boyle WA, Cheng TL, Clerc J, Fuller NW, Gerson AR, Jonasson KA, Rogers EJ *et al.* (2018) Common condition indices are no more effective than body mass for estimating fat stores in insectivorous bats. *J Mammal* 99: 1065–1071.
- McMichael L, Edson D, Smith C, Mayer D, Smith I, Kopp S, Meers J, Field H (2017) Physiological stress and Hendra virus in flying-foxes (*Pteropus* spp.), Australia. *PLoS One* 12. [10.1371/journal.pone.0182171](https://doi.org/10.1371/journal.pone.0182171).
- Measor KR, Leavell BC, Brewton DH, Rumschlag J, Barber JR, Razak KA (2017) Matched behavioral and neural adaptations for low sound level echolocation in a gleaning bat, *Antrozous pallidus*. *eNeuro* 4: 0018–0017.2017.
- Medellin RA, Rivero M, Ibarra A, De La Torre JA, Gonzalez-Terrazas TP, Torres-Knoop L, Tschapka M (2018) Follow me: foraging distances of *Leptonycteris yerbabuenae* (Chiroptera: Phyllostomidae) in Sonora determined by fluorescent powder. *J Mammal* 99: 306–311.
- Meyer JS, Novak MA (2012) Minireview: hair cortisol: a novel biomarker of hypothalamic-pituitary-adrenocortical activity. *Endocrinology* 153: 4120–4127.
- Michonneau F, Brown JW, Winter DJ (2016) rotl: an R package to interact with the Open Tree of Life data. *Methods Ecol Evol* 7: 1476–1481.
- Mickleburgh SP, Hutson AM, Racey PA (2002) A review of the global conservation status of bats. *Oryx* 36: 18–34.
- Montillo M, Caslini C, Peric T, Prandi A, Netto P, Tubaro F, Pedrotti L, Bianchi A, Mattiello S (2019) Analysis of 19 minerals and cortisol in red deer hair in two different areas of the stelvio national park: a preliminary study. *Animals* 9: 492.
- Myhrvold NP, Baldrige E, Chan B, Sivam D, Freeman DL, Ernest SKM (2015) An amniote life-history database to perform comparative analyses with birds, mammals, and reptiles. *Ecology* 96: 3109–3000.
- Norberg UM, Rayner JMV (1987) Ecological morphology and flight in bats (Mammalia; Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation. *Philos Trans R Soc London B, Biol Sci* 316: 335–427.
- Norris DO, Carr JA (2013) An overview of chemical bioregulation in vertebrates. In *Vertebrate Endocrinology*. Elsevier, pp. 1–22
- O'Donnell CFJ (2000) Conservation status and causes of decline of the threatened New Zealand long-tailed bat *Chalinolobus tuberculatus* (Chiroptera: Vespertilionidae). *Mamm Rev* 30: 89–106.
- Oelbaum PJ, Fenton MB, Simmons NB, Broders HG (2019) Community structure of a Neotropical bat fauna as revealed by stable isotope analysis: not all species fit neatly into predicted guilds. *Biotropica* 51: 719–730.
- Orona Castillo I, Sangerman-Jarquín DM, Fortis Hernández M, Vázquez Vázquez C, Gallegos Robles MÁ (2018) Producción y comercialización de nuez pecanera (*Carya illinoensis* Koch) en el norte de Coahuila, México. *Rev Mex Ciencias Agrícolas* 4: 461–476.
- Ouyang JQ, Sharp PJ, Dawson A, Quetting M, Hau M (2011) Hormone levels predict individual differences in reproductive success in a passerine bird. *Proc R Soc B Biol Sci* 278: 2537–2545.
- Pagel M (1999) Inferring the historical patterns of biological evolution. *Nature* 401: 877–884.
- Paradis E, Claude J, Strimmer K (2004) APE: Analyses of Phylogenetics and Evolution in R language. *Bioinform Appl NOTE* 20: 289–290.
- Patterson BD, Dick CW, Dittmar K (2007) Roosting habits of bats affect their parasitism by bat flies (Diptera: Streblidae). *J Trop Ecol* 23: 177–189.
- Patterson SH, Hahn TP, Cornelius JM, Breuner CW (2014) Natural selection and glucocorticoid physiology. *J Evol Biol* 27: 259–274.

- Pavan AC, Marroig G (2016) Integrating multiple evidences in taxonomy: species diversity and phylogeny of mustached bats (Mormoopidae: Pteronotus). *Mol Phylogenet Evol* 103: 184–198.
- Pearson Murphy BE (2007) *Glucocorticoids, Overview*. Encyclopedia of Stress. Elsevier Inc., pp. 198–210
- Pennell MW (2015) *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology: Concepts and Practice*. Edited by László Zsolt Garamszegi Vol 64, pp. 161–163 Systematic Biology
- Plowright RK, Field HE, Smith C, Divljan A, Palmer C, Tabor G, Daszak P, Foley JE (2008) Reproduction and nutritional stress are risk factors for Hendra virus infection in little red flying foxes (*Pteropus scapulatus*). *Proc R Soc B Biol Sci* 275: 861–869.
- Postawa T, Szubert-Kruszyńska A (2014) Is parasite load dependent on host aggregation size? The case of the greater mouse-eared bat *Myotis myotis* (Mammalia: Chiroptera) and its parasitic mite *Spinturnix myoti* (Acari: Gamasida). *Parasitol Res* 113: 1803–1811.
- Pragst F, Balikova MA (2006) State of the art in hair analysis for detection of drug and alcohol abuse. *Clin Chim Acta* 370: 17–49.
- Racey P (2009) Reproductive assessment in bats. In T Kunz, ed, *Ecological and Behavioral Methods for the Study of Bats*, EdEd 2. Smithsonian Press, Washington, DC, pp. 31–45
- Racey PA (2013) Bat Conservation: Past, Present and Future. In *Bat Evolution, Ecology, and Conservation*. Springer New York, New York, NY, pp. 517–532
- Rakotoniaina JH, Kappeler PM, Kaesler E, Hämäläinen AM, Kirschbaum C, Kraus C (2017) Hair cortisol concentrations correlate negatively with survival in a wild primate population. *BMC Ecol* 17: 1–13.
- Reeder DAM, Kosteczko NS, Kunz TH, Widmaier EP (2004) Changes in baseline and stress-induced glucocorticoid levels during the active period in free-ranging male and female little brown myotis, *Myotis lucifugus* (Chiroptera: Vespertilionidae). *Gen Comp Endocrinol* 136: 260–269.
- Reeder DM, Kramer KM (2005) Stress in free-ranging mammals: integrating physiology, ecology, and natural history. *J Mammal* 86: 225–235.
- Reid F (2009) *A Field Guide to the Mammals of Central America & Southeast Mexico*, EdEd 2. Oxford University Press, New York, Oxford
- Ricklefs RE, Wikelski M (2002) The physiology/life-history nexus. *Trends Ecol Evol* 17: 462–468.
- Romero LM (2002) Seasonal changes in plasma glucocorticoid concentrations in free-living vertebrates. *Gen Comp Endocrinol* 128: 1–24.
- Romero LM (2004) Physiological stress in ecology: Lessons from biomedical research. *Trends Ecol Evol* 19: 249–255.
- Russell E, Koren G, Rieder M, Van Uum S (2012) Hair cortisol as a biological marker of chronic stress: current status, future directions and unanswered questions. *Psychoneuroendocrinology* 37: 589–601.
- Santana SE, Dial TO, Eiting TP, Alfaro ME (2011) Roosting ecology and the evolution of pelage markings in bats. *PLoS One* 6: e25845.
- Sapolsky RM, Romero LM, Munck AU (2000) How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 21: 55–89.
- Schneeberger K, Cziráj G, Voigt CC (2013) Measures of the constitutive immune system are linked to diet and roosting habits of neotropical bats. *PLoS One* 8:1. [10.1371/journal.pone.0054023](https://doi.org/10.1371/journal.pone.0054023).
- Schulte PM (2014) What is environmental stress? Insights from fish living in a variable environment. *J Exp Biol* 217: 23–34.
- Sharpley CF, McFarlane JR, Slominski A (2012) Stress-linked cortisol concentrations in hair: what we know and what we need to know. *Rev Neurosci* 23: 111–121.
- Sheriff MJ, Dantzer B, Delehanty B, Palme R, Boonstra R (2011) Measuring stress in wildlife: techniques for quantifying glucocorticoids. *Oecologia* 166: 869–887.
- Sikes RS, Bryan JA (2016) 2016 Guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *J Mammal* 97: 663–688.
- Simmons NB, Cirranello AL (2020) Bat Species of the World: a taxonomic and geographic database batnames |home. <https://batnames.org/> (last accessed 22 June 2020).
- Speakman J, Thomas D (2003) Physiological ecology and energetics of bats. In T Kunz, M Fenton, eds, *Bat Ecology*. The University of Chicago Press, Chicago, pp. 430–490
- Speakman JR (1997) *Doubly Labeled Water: Theory and Practice*. Chapman & Hall Ltd, London, UK
- Stahlschmidt P, Brühl CA (2012) Bats as bioindicators—the need of a standardized method for acoustic bat activity surveys. *Methods Ecol Evol* 3: 503–508.
- Stearns SC (1992) *The Evolution of Life Histories*. Oxford University, Oxford
- Sullivan AR, Bump JK, Kruger LA, Peterson RO (2012) Bat-cave catchment areas: using stable isotopes ( $\delta D$ ) to determine the probable origins of hibernating bats. *Ecol Appl* 22: 1428–1434.
- Téllez HLA, Iñiguez-Dávalos LI, Olvera-Vargas M, Vargas-Contreras JA, Herrera-Lizaola OA (2018) Bats associated to caves in Jalisco, Mexico. *Therya* 9: 29–40.
- Terwissen CV, Mastromonaco GF, Murray DL (2013) Influence of adrenocorticotrophin hormone challenge and external factors (age, sex, and body region) on hair cortisol concentration in Canada lynx (*Lynx canadensis*). *Gen Comp Endocrinol* 194: 162–167.
- Tilbrook A (2000) Effects of stress on reproduction in non-rodent mammals: the role of glucocorticoids and sex differences. *Rev Reprod* 5: 105–113.
- Tiunov M, Makarikova T (2007) Seasonal molting in *Myotis petax* (Chiroptera) in the Russian Far East. *Acta Chiropt* 9: 538–538.
- Torres-Flores JW, López-Wilchis R, Soto-Castruita A (2012) Population dynamics, roost selection and reproductive patterns of some cave bats from Western of Mexico. *Rev Biol Trop* 60: 1369–1389.



- Voigt CC, Kingston T (2015) Bats in the anthropocene. In *Bats in the Anthropocene: Conservation of Bats in a Changing World*. Springer International Publishing, Cham, Switzerland, pp. 1–9
- Voigt CC, Voigt-Heucke SL, Schneeberger K (2012) Isotopic data do not support food sharing within large networks of female vampire bats (*Desmodus rotundus*). *Ethology* 118: 260–268.
- Wada H (2008) Glucocorticoids: mediators of vertebrate ontogenetic transitions. *Gen Comp Endocrinol* 156: 441–453.
- Wada H, Yates DE, Evers DC, Taylor RJ, Hopkins WA (2010) Tissue mercury concentrations and adrenocortical responses of female big brown bats (*Eptesicus fuscus*) near a contaminated river. *Ecotoxicology* 19: 1277–1284.
- Wikelski M, Cooke SJ (2006) Conservation physiology. *Trends Ecol Evol* 21: 38–46.
- Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W (2014) EltonTraits 1.0: species-level foraging attributes of the world's birds and mammals. *Ecology* 95: 2027–2027.
- Wingfield JC, Romero LM (2011) Adrenocortical responses to stress and their modulation in free-living vertebrates. In *Comprehensive Physiology*. John Wiley & Sons, Inc., Hoboken, NJ, USA, pp. 211–234
- Wingfield JC, Sapolsky RM (2003) Reproduction and resistance to stress: when and how. *J Neuroendocrinol* 15: 711–724.
- Zuloaga DG, Heck AL, De Guzman RM, Handa RJ (2020) Roles for androgens in mediating the sex differences of neuroendocrine and behavioral stress responses. *Biol Sex Differ* 11: 1–18.